

## Lost water and nitrogen resources due to EU consumer food waste

This content has been downloaded from IOPscience. Please scroll down to see the full text.

2015 Environ. Res. Lett. 10 084008

(<http://iopscience.iop.org/1748-9326/10/8/084008>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 84.123.192.126

This content was downloaded on 12/08/2015 at 14:46

Please note that [terms and conditions apply](#).

## Environmental Research Letters



## LETTER

## Lost water and nitrogen resources due to EU consumer food waste

## OPEN ACCESS

RECEIVED  
24 February 2015

REVISED  
11 June 2015

ACCEPTED FOR PUBLICATION  
10 July 2015

PUBLISHED  
12 August 2015

Content from this work may be used under the terms of the [Creative Commons Attribution 3.0 licence](#).

Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.



D Vanham, F Bouraoui, A Leip, B Grizzetti and G Bidoglio

European Commission, Joint Research Centre, Institute for Environment and Sustainability, Via E. Fermi 2749, I-21027 Ispra (VA), Italy

E-mail: [davy.vanham@yahoo.de](mailto:davy.vanham@yahoo.de)

**Keywords:** food waste, consumer, EU, water footprint, nitrogen footprint, uncertainty, footprint

**Abstract**

The European Parliament recently called for urgent measures to halve food waste in the EU, where consumers are responsible for a major part of total waste along the food supply chain. Due to a lack of data on national food waste statistics, uncertainty in (consumer) waste quantities (and the resulting associated quantities of natural resources) is very high, but has never been previously assessed in studies for the EU. Here we quantify: (1) EU consumer food waste, and (2) associated natural resources required for its production, in term of water and nitrogen, as well as estimating the uncertainty of these values. Total EU consumer food waste averages 123 (min 55–max 190) kg/capita annually (kg/cap/yr), i.e. 16% (min 7–max 24%) of all food reaching consumers. Almost 80%, i.e. 97 (min 45–max 153) kg/cap/yr is avoidable food waste, which is edible food not consumed. We have calculated the water and nitrogen (N) resources associated with avoidable food waste. The associated blue water footprint (WF) (the consumption of surface and groundwater resources) averages 27 litre per capita per day (min 13–max 40 l/cap/d), which slightly exceeds the total blue consumptive EU municipal water use. The associated green WF (consumptive rainwater use) is 294 (min 127–max 449) l/cap/d, equivalent to the total green consumptive water use for crop production in Spain. The nitrogen (N) contained in avoidable food waste averages 0.68 (min 0.29–max 1.08) kg/cap/yr. The food production N footprint (any remaining N used in the food production process) averages 2.74 (min 1.02–max 4.65) kg/cap/yr, equivalent to the use of mineral fertiliser by the UK and Germany combined. Among all the food product groups wasted, meat accounts for the highest amounts of water and N resources, followed by wasted cereals. The results of this study provide essential insights and information on sustainable consumption and resource efficiency for both EU policies and EU consumers.

**1. Introduction**

Global food security will be one of mankind's main challenges this century. A key question is whether humanity has the natural resource base to feed itself equitably and sustainably both now and in the future. Currently, 842 million people are chronically undernourished [1], while 1.46 billion adults are overweight, of whom 502 million are obese [2]. Within the EU, 227 million (58% of all) adults are overweight, of whom 91 million (23% of all adults) are obese [3, 4]. By 2050, a projected 9–10 billion people should be provided with a healthy diet in a sustainable way [5]. In creating a more sustainable food system, solutions need to combine demand-side and supply-side options [6].

Demand-side solutions should target dietary habits as well as food waste [7–9]. This paper focuses on food waste.

According to the UN's Food and Agriculture Organization (FAO) [8], roughly one-third of food produced for human nutrition is either lost or wasted globally, which amounts to approximately 1.3 billion tonnes per year. Food is lost or wasted throughout the entire supply chain, from initial agricultural production down to final household consumption. In industrial countries, the fraction of food wasted by consumers of the total food lost and wasted is high. According to this FAO study [8], the food wasted per-capita by consumers in Europe and North America is about 95 to 115 kg yr<sup>-1</sup>. Total per-capita food loss and

waste along the entire supply chain in Europe and North America is 280–300 kg yr<sup>-1</sup>. Thus, consumer food waste represents about one-third of this amount. A study on consumer food waste in the EU-27 [10] quantified total food waste per capita at 101 kg yr<sup>-1</sup> (76 kg yr<sup>-1</sup> for households and 25 kg yr<sup>-1</sup> for the food service/catering sector).

An essential limitation of previous quantification studies on consumer food waste for Europe—e.g. [8] or the EU—e.g. [10, 11]—is that they only use one value (total and/or per product/product group), thereby not taking into account the uncertainty and lack of reliable data on food waste statistics. The FAO study [8] on food losses and waste only uses UK data for consumer food waste in Europe [12]. The study on food waste carried out on behalf of the European Commission [10] quantifies total consumer food waste for each EU Member State based on assembled statistical data, but gives no range on uncertainty and does not differentiate between product groups. A recent study on behalf the European Parliament [13] acknowledges that the estimations in the latter study are conservative, and probably underestimate.

A number of studies have quantified particular natural resources (including energy [14]) associated with food waste, often by means of environmental footprints, the latter being an umbrella term for the different footprint concepts that have been developed during the last two decades [15]. Well-established footprint indicators include the water footprint (WF) [16], nitrogen footprint (NF) [11, 17, 18], land footprint (LF) [19] and carbon footprint (CF) [20]. The WF is an indicator of direct and indirect water use. The concept has been introduced into water-management science to show the importance of consumption patterns and the global dimensions of good water governance [16]. The NF comprises the sum of the nitrogen contained in food (based upon protein content) and the food production NF. The LF measures the appropriation of land as a resource. The CF measures the emission of greenhouse gases (GHGs) into the atmosphere. For example, Chapagain and James [21] calculated the WF and CF of household food and drink waste in the UK. The FAO [22] quantified the CF, WF and LF of global food wastage (i.e. both food loss and food waste). Kummu *et al* [23] quantified the WF as well as the cropland and fertiliser use associated with food wasted on a global level.

The aim of this study was to quantify EU consumer food waste, taking account of uncertainty. In particular, we have quantified not only the total amount of food waste but also the food waste per product group. We have also differentiated between total food waste and avoidable food waste. In addition, we have quantified the associated natural resources required for the production of this food waste, in terms of water and nitrogen. For this, we have used the WF and NF concepts. We chose these two footprints as they are well established and because detailed studies

have been made on the WF and NF of EU food production and consumption [9, 24–26]. Future research should include additional resources/footprints.

## 2. Methodology

### 2.1. General

We have quantified consumer food waste for the EU-28 with the associated uncertainty. We have analysed average annual values for the period 1996–2005. During this period, the average population of the EU was 487 million. The methodology is presented below and an overview of the workflow of the methodology is given in figure 1, with explanations of the abbreviations used throughout the manuscript provided in table 1.

### 2.2. Quantification of food consumption data (FOOD\_CONS)

As a first step, average EU food consumption data have been quantified by taking food supply quantity data (FOOD\_SUPPLY) of different product groups from the FAO Food balance sheets (FBS) [27]. These provide data on the amount of food that reaches the consumer in private households, as well as that in the non-household sector, i.e. catering establishments, boarding schools, hospitals, prisons, armed forces' bases and other communities. The data are given on an 'as purchased' basis, i.e. as the food leaves the retail shop or enters the household by other means.

Quantities are provided on the basis of 'primary equivalents'. Within the FAO FBS, food data are standardized in that processed commodities are converted back to their 'primary equivalent'. This is for standardization (different countries report their data to the FAO), simplification and limitation of the number of commodities within the FBS. The latter implies that it is beneficial to reduce the amount of data, and therefore the number of commodities involved, to a level and size more suited to analytical purposes. E.g., instead of listing flour of wheat, bread or pasta separately in the FBS, they are quantified as wheat equivalent. Similarly, meat (reaching consumers in many forms, e.g. for chicken as—amongst others—a whole chicken, chicken filet, sausages or chicken nuggets) is quantified as carcass weight in the FBS.

We have estimated food consumption at the point where it reaches the consumer, using a first correction factor (CORR1) which accounts for product primary equivalent conversion (equation (1)). To calculate total and avoidable wasted food quantities from FOOD\_SUPPLY, two correction factors are applied. The first factor (CORR1) accounts for product primary equivalent conversions (equation (1)) and the second for food waste (CORR2).

The specific values for CORR1 have been taken from [28]. For example, the CORR1 of wheat equivalent is 0.8, as only 80% of the original wheat weight




Figure 1. Workflow of the methodology; the equations are listed in table 2.

Table 1. Abbreviations used in the letter.

Abbreviation	Definition
EU	European Union
WF	Water footprint
$WF_{prod}$	Water footprint of production
$WF_{cons}$	Water footprint of consumption
$WF_{cons, agr}$	Agricultural water footprint of production
$WF_{cons, agr, gn}; WF_{cons, agr, bl}$	Green; blue agricultural water footprint of consumption
NF	Nitrogen footprint
$N_{cons}$	Nitrogen contained in food
$NF_{prod}$	Food production NF
FAO	Food and Agriculture Organization of the United Nations
FBS	Food balance sheet
FOOD_SUPPLY	Food supply quantity (data)
FOOD_CONS	Food consumption quantity (data)
CORR1	Correction factor 1, which accounts for product primary equivalent conversions
CORR2	Correction factor 2, which accounts for food waste
CORR2_TOT	Correction factor 2, which accounts for total food waste
CORR2_AV	Correction factor 2, which accounts for avoidable food waste
CORR2_PA	Correction factor 2, which accounts for potentially avoidable food waste
CORR2_NA	Correction factor 2, which accounts for non-avoidable food waste
WASTE_TOT	Total food waste quantity
WASTE_AV	Avoidable food waste quantity
WASTE-NA	Non-avoidable food waste quantity
DEFRA	UK Department for Environment, Food & Rural Affairs
Mt	Million tonnes
l/cap/d	Litre per capita per day

**Table 2.** Equations used to compute waste amounts (equations (1)–(6)) and related resources (equations (7)–(9)).

FOOD_SUPPLY × CORR1 =	FOOD_CONS	(1)
with: FOOD_SUPPLY =	Food supply quantity as given in FAO Food Balance Sheets (FBS). Food supply is food (in tonnes) reaching the consumer, i.e. in private households, as well as the food service/catering sector	
	CORR1 =	Correction factor 1, which accounts for product primary equivalent conversions
	FOOD_CONS =	Food consumption.
FOOD_CONS × CORR2_TOT =	WASTE_TOT	(2)
FOOD_CONS × CORR2_AV =	WASTE_AV	(3)
FOOD_CONS × CORR2_NA =	WASTE_NA	(4)
With: CORR2 =	Correction factor 2, which accounts for food waste	
	CORR2_TOT =	Correction factor 2, which accounts for total food waste
	CORR2_AV =	Correction factor 2, which accounts for avoidable food waste
	CORR2_NA =	Correction factor 2, which accounts for non-avoidable food waste
	WASTE_TOT =	Amount of total consumer food waste
	WASTE_AV =	Amount of avoidable consumer food waste
	WASTE_NA =	Amount of non-avoidable consumer food waste
WASTE_TOT =	WASTE_AV + WASTE_NA	(5)
FOOD_CONS =	INTAKE + WASTE_TOT	(6)
With: INTAKE =	Amount of food actually consumed	
WASTE_AV <sub>CORR</sub> =	$\frac{WASTE\_AV}{1 - CORR2\_NA}$	(7)
with	WASTE_AV <sub>CORR</sub> =	Corrected amount of avoidable consumer food waste to relate to the resource footprint (WF or NF) of WASTE_AV
	CORR2_NA =	$\frac{WASTE\_NA}{FOOD\_CONS}$ as obtained from (1) and (4):
WF <sub>cons,agr</sub> of WASTE_AV =	$\left(\frac{WASTE\_AV_{CORR}}{FOOD\_CONS}\right) \times WF_{cons,agr}$	(8)
with	WF <sub>cons,agr</sub> =	WF <sub>cons,agr</sub> of FOOD_CONS
NF of WASTE_AV =	$\left(\frac{WASTE\_AV_{CORR}}{FOOD\_CONS}\right) \times NF$	(9)
With	NF =	NF of FOOD_CONS

remains after conversion to wheat flour. In addition to the first factor values listed in the latter publication, 7.4 milk equivalent has been taken for cheese, as obtained from [3], 7 milk equivalent for cream, and 1 milk equivalent for yoghurt. After applying CORR1, food consumption data (FOOD\_CONS) have been quantified.

### 2.3. Quantification of food waste amounts

Total and avoidable quantities of wasted food from FOOD\_SUPPLY have been obtained by applying a second correction factor for food waste (CORR2) on food consumption quantities (FOOD\_CONS). This factor relates either to total food waste (CORR2\_TOT) or to avoidable food waste (CORR2\_AV), as we have differentiated between total (WASTE\_TOT) and avoidable (WASTE\_AV) food waste. The latter excludes non-avoidable food waste (WASTE\_NA) such as meat bones, egg shells, fruit stones or the peel of certain fruit and vegetables. Avoidable food waste means food that at some point prior to disposal was edible. The relevant equations (2)–(6) are listed in table 2 below.

Based upon FOOD\_CONS, total (WASTE\_TOT, equation (2)) and avoidable food (WASTE\_AV, equation (3)) waste quantities are thus calculated by means of CORR2\_TOT and CORR2\_AV. Non-avoidable food waste (WASTE\_NA) is the difference between WASTE\_TOT and WASTE\_AV (equation (5)).

CORR2\_TOT and CORR2\_AV values have been obtained for the product groups from/based upon six selected national studies, i.e.

- the UK [29],
- the Netherlands [30],
- Denmark [31],
- Finland [32],
- Germany [33],
- Romania [34].

For other EU countries, either no data were found or they were unreliable. These values are shown in table 3 below.

Only estimations of household waste have been quantified in these studies. For the non-household (food service/catering) sector part of FOOD\_CONS, the same CORR2 values have been taken.

The most detailed assessment was done in the UK [29] for the year 2007 (a field study of over 2000 households), where a distinction is made between avoidable waste, potentially avoidable waste and unavoidable waste. Where purchased amounts of specific products or product groups are missing, DEFRA (the UK's Department for Environment, Food & Rural Affairs) statistics have been used. Apart from avoidable and non-avoidable food waste, the UK study also

**Table 3.** Estimates of food waste fractions (CORR2 in %) of household food purchases (FOOD\_CONS) by food group, based upon literature values.

Product group	UK [29]				The Netherlands, based upon [30]			Denmark, based upon [31]			Finland [32]	Germany, based upon [33]		Romania [34]
	CORR2_AV	CORR2_PA	CORR2_NA	CORR2_TOT	CORR2_AV	CORR2_NA	CORR2_TOT	CORR2_AV	CORR2_NA	CORR2_TOT	CORR2_AV	CORR2_AV	CORR2_TOT	CORR2_AV
Cereals	26	3	0	29	12	0	12	12	1	13	13	11–15		>0 and <10
Potatoes	17	28	0	45	5	4	9				19			
Vegetables	23	13	9	45	9	11	20	5	11	16	19	15–20	18–23	>0 and <10
Fruit	19	3	19	41	9	16	25	5	11	16	13	8–10	18–23	>0 and <10
Sugar														
Pulses, nuts and oil crops														
Crop oils	4	12 <sup>a</sup>	1	17										
Animal fats	4	12 <sup>a</sup>	1	17										
Stimulants														
Spices														
Alcoholic beverages	5	0	0	5										
Meat	11	3	9	23	6	2	8	5	2	7	7	4–6	11–15	>0 and <10
Fish	11	3	9	23	2	7	9	5	2	7	7	4–6	11–15	>0 and <10
Eggs	5	0	12	17	2	6	8				7			
Milk	7	0	0	7	5	0	5				17	3–4	7–9	>0 and <10
Yoghurt	13	0	0	13	5	0	5				17	3–4	7–9	>0 and <10
Cheese	10	0	0	10	3	1	4				17	3–4	7–9	>0 and <10

<sup>a</sup> In ([29]) the potentially avoidable waste for oils and fats includes all oils and fats which have only been used once—however, this is not recommended in other countries, therefore a second use is discarded in this analysis and the potentially avoidable waste proportion reduced to 0%.



**Table 4.** Statistical normal distribution values (av = average; stdev = standard deviation; max = maximum; min = minimum) of total and avoidable food waste fractions (CORR2\_TOT and CORR2\_AV in %), based upon the literature values in table 3. For sugar, stimulants, spices and alcoholic beverages, correction factors are assumed to be uniformly distributed.

Product group	Total waste				Avoidable waste			
	av	stdev	max	min	av	stdev	max	min
Cereals	17.12	8.6	29	5	17.12	8.6	29	5
Potatoes	25.5	14.2	45	5	25.2	15	45	5
Sugar			10	5			10	5
Pulses	5	1.8	7	2	4.7	1.9	7	2
Crop oils	5	1.8	7	2	4.7	1.9	7	2
Vegetables	26.2	13.9	45	5	20.9	11.5	36	4
Fruit	25.5	12	41	5	12.6	6.8	22	4
Stimulants			10	5			10	5
Spices			10	5			10	5
Alcoholic beverages			7	3			7	3
Meat	14.5	6.6	23	4	7.7	4.6	14	2
Animal fats	5	1.8	7	2	5	1.8	7	2
Eggs	11.9	3.9	17	6	5.1	1.8	7	2
Milk (and yoghurt)	7	2.8	17	2	5.2	3	17	2
Cheese	7.9	3.2	17	2	6	3.9	17	2
Cream	5.2	3	17	2	5.2	3	17	2
Fish	14.5	6.5	23	4	7.4	4.8	14	2

identifies potentially avoidable food waste. However, in our study we have added this fraction (CORR2\_PA) to avoidable food waste (CORR2\_AV) for the UK statistics, as other national studies do not account for potentially avoidable food waste.

Table 3 shows that for some product groups detailed statistics are available for several countries, e.g. for cereals, vegetables, fruit, meat, milk and milk products. For other product groups, statistics are only available for selected countries (e.g. alcoholic beverages or crop oils) and for some product groups, no statistics are available (e.g. sugar or stimulants). The data compiled in table 3 show that for all product groups: (1) CORR2\_TOT and CORR2\_AV values are highest in the UK (except for milk/milk products where Finland has a higher value); and (2) CORR2\_AV values are amongst the lowest in Romania. To account for any uncertainty in the estimations, we have assumed that Romania represents the minimum CORR2 values and the UK the maximum values for EU countries, and that all other values are between these two extremes. Schneider [35] has already pointed out that in Europe, the proportion of household expenditure on food is highest in Eastern Europe (e.g. 44.2% in Romania) and that this might be expected to have a significant impact on food waste behaviour. The UK has already been identified as the EU country with the highest per capita consumer food waste [10].

To compute the CORR2\_TOT and CORR2\_AV distributions for the EU, based upon the six selected national studies, we have made some assumptions. In particular, we have assumed that the correction factors (CORR2\_TOT and CORR2\_AV) per product group to calculate WASTE\_TOT and WASTE\_AV are distributed normally. Their means and standard

deviations (as shown in table 4 below) were calculated from the available country data (table 3). To calculate these mean EU values from the six national values, these national values are weighted according to national populations. For example, the values for Germany are weighted at 43% and those of the Netherlands at 8%. For sugar, stimulants, spices and alcoholic beverages the associated correction factors were assumed to be uniformly distributed, as detailed data are missing.

#### 2.4. Quantification of related water resources (the water footprint or WF)

In our study, we used the WF concept. For the purpose of water resources management (e.g. in the EU, as in our study), a geographical WF assessment is relevant. It is important to distinguish between the WF of production ( $WF_{prod}$ ) and the WF of consumption ( $WF_{cons}$ ) of a region [16]. The first refers to the total use of domestic water resources within the region (for producing goods and services for domestic consumption or export). The second refers to the use of domestic and foreign water resources behind all the goods and services consumed domestically.

Dependent on the authors, a WF comprises either three (blue, green and grey) or two (blue and green) components. As defined in [36], green water is the soil water retained in the unsaturated zone, which is formed by precipitation and is available to plants; blue water refers to water in rivers, lakes, wetlands and aquifers. Irrigated agriculture receives blue water (from irrigation) as well as green water (from precipitation), while rain-fed agriculture only receives green water. Thus, the green WF is the rainwater consumed by crops. The grey WF is an indicator of the

**Table 5.** Average annual (1996–2005) FOOD\_SUPPLY and FOOD\_CONS values (in kg/cap/yr) of different product groups for the EU. The  $WF_{\text{cons,agr,gn}}$  and  $WF_{\text{cons,agr,bl}}$  (in l/cap/d) and the  $N_{\text{cons}}$  and  $NF_{\text{prod}}$  (in kg/cap/yr) are also shown.

Product group	FOOD_SUPPLY (kg/cap/yr)	FOOD_CONS (kg/cap/yr)	Green $WF_{\text{cons,agr}}$ (l/cap/d)	Blue $WF_{\text{cons,agr}}$ (l/cap/d)	$N_{\text{cons}}$ (kg/cap/yr)	$NF_{\text{prod}}$ (kg/cap/yr)
Cereals	123.7	93.3	337	21	1.7	1.9
Potatoes	81.9	81.9	33	6	0.2	0.3
Sugar	39.7	39.7	158	30	0.0	0.3
Pulses, nuts and oil crops	10.3	10.3	95	10	0.2	0.1
Crop oils	18.5	18.5	293	23	0.0	0.4
Vegetables	120.8	120.8	47	14	0.2	0.3
Fruit	98.8	98.8	115	30	0.1	0.3
Stimulants	7.5	7.5	322	2	0.1	0.0
Spices	0.5	0.5	9	1	0.0	0.0
Alcoholic beverages	109.3	109.3	84	6	0.1	1.2
Meat	88.5	59.4	1247	82	1.7	18.3
Animal fats	13.3	13.3	103	6	0.0	0.5
Eggs	12.4	12.4	68	4	0.2	0.6
Milk (excluding butter) <sup>a</sup>	234.1 (including cheese, cream and yoghurt)	95.3 (milk and yoghurt—yoghurt 1 l milk for 1 kg)	192	15	0.5	2.6
Cheese <sup>a</sup>	0	15.2 (112.5 milk eq.—7.4 l milk for 1 kg)	227	17	0.6	3.1
Cream <sup>a</sup>	0	3.8 (26.3 milk eq.—7.0 l milk for 1 kg)	53	4	0.1	0.7
Fish and seafood	21.2	8.5	0	0	0.4	1.8
Total	980.5	788.5	3383	270	6.0	32.4

<sup>a</sup> FOOD\_CONS values for cheese and cream are taken from FAOSTAT food supply statistics. Milk equivalent (milk eq.) values are computed from EUROSTAT-data.

degree of water pollution [37]. However, many authors perceive this component critically [24, 38, 39].

To quantify the water resources related to these wasted food amounts, we have used the WF of consumption ( $WF_{\text{cons}}$ ) concept [16, 37]. In particular, we have assessed the green and blue  $WF_{\text{cons}}$  of agricultural products ( $WF_{\text{cons,agr}}$ ) for wasted food. Therefore, the  $WF_{\text{cons,agr}}$  refers to the use of domestic and foreign water resources for all agricultural goods that are consumed domestically. The green  $WF_{\text{cons,agr}}$  ( $WF_{\text{cons,agr,gn}}$ ) represents the consumptive rainwater use for the production of crops and animal products which EU citizens consume. Similarly, the blue  $WF_{\text{cons,agr}}$  ( $WF_{\text{cons,agr,bl}}$ ) represents the consumptive use of water from rivers, lakes and groundwater for this production.

As a first step, the  $WF_{\text{cons,agr}}$  of FOOD\_CONS needs to be quantified. To assess the  $WF_{\text{cons,agr}}$  we have followed the methodology of the Global Water Footprint Standard developed by the Water Footprint Network [36]. National green and blue  $WF_{\text{cons,agr}}$  data are accessed from [16, 40] and aggregated to product groups at the EU level. As a result, the blue and green  $WF_{\text{cons,agr}}$  ( $WF_{\text{cons,agr,gn}}$  and  $WF_{\text{cons,agr,bl}}$ ) of FOOD\_CONS are obtained.

As a next step, the  $WF_{\text{cons,agr}}$  ( $WF_{\text{cons,agr,gn}}$ ) and ( $WF_{\text{cons,agr,bl}}$ ) of avoidable waste amounts (WASTE\_AV) needs to be quantified. To do this, first the

unavoidable fraction (CORR2\_NA) must be added to WASTE\_AV because the  $WF_{\text{cons,agr}}$  of FOOD\_CONS relates to the whole product. Thus we have calculated a new component WASTE\_AV<sub>CORR</sub> by means of equation (7) (table 2).

When both the  $WF_{\text{cons,agr}}$  of FOOD\_CONS and WASTE\_AV<sub>CORR</sub> are calculated, the  $WF_{\text{cons,agr,gn}}$  and  $WF_{\text{cons,agr,bl}}$  for WASTE\_AV are computed using equation (8) (table 2).

To clarify, we present the example of vegetables. Using equation (1), we have calculated the FOOD\_CONS value for vegetables as 120.8 kg/cap/yr (table 5). The related  $WF_{\text{cons,agr,gn}}$  of this amount is 47 l/cap/d (table 5). With an average avoidable food waste fraction (CORR2\_AV) of 20.9% (table 4), we have used equation (3) to calculate the average amount of avoidable consumer fruit waste (WASTE\_AV) as 25.3 kg/cap/yr.

In order to calculate the  $WF_{\text{cons,agr,gn}}$  of this avoidable food waste, first we added the unavoidable fraction of the amount of waste. For example, when an onion is wasted the avoidable waste quantity (WASTE\_AV) is the onion without its skin. The  $WF_{\text{cons,agr,gn}}$  of all onions consumed and wasted by EU consumers (FOOD\_CONS of onions) relates to the whole product, including the skin. Therefore the skin must be added to the wasted onion to calculate its  $WF_{\text{cons,agr,gn}}$ .



For all vegetables, the corrected avoidable waste amount ( $WASTE\_AV_{CORR}$ )—which is only useful to calculate the related footprint—is computed by means of equation (7).

$$WASTE\_AV_{CORR} = \frac{WASTE\_AV}{1 - CORR2\_NA} \quad (\text{equation (7)}) = \frac{25.3}{1 - 0.053} = 26.7 \text{ kg/cap/yr}$$

$$\text{with } CORR2\_NA = \frac{WASTE\_NA}{FOOD\_CONS} = \frac{6.4}{120.8} = 0.053$$

$$\text{with } WASTE\_NA \text{ (equation (4))} = FOOD\_CONS \times CORR2\_NA = 120.8 \text{ kg/cap/yr} \times 5.3\% = 6.4 \text{ kg/cap/yr}$$

The  $WF_{cons,agr,gn}$  of avoidable vegetable waste is then calculated using equation (8):

$$\begin{aligned} & \left( \frac{WASTE\_AV_{CORR}}{FOOD\_CONS} \right) \times WF_{cons,agr} \\ & = \left( \frac{26.7}{120.8} \right) \times 47 \text{ l/cap/d} = 10.4 \text{ l/cap/d.} \end{aligned}$$

## 2.5. Quantification of related nitrogen resources (the nitrogen footprint or NF)

To quantify the nitrogen resources related to these wasted food amounts, we have used the NF concept [11, 18, 26]. In particular, we have assessed the nitrogen contained in food ( $N_{cons}$ ) and in food production NF ( $NF_{prod}$ ) for avoidable food waste. The  $N_{cons}$  quantifies the N-content within a product (based upon the protein content). The  $NF_{prod}$  is any nitrogen that has been used in the food chain and has been lost to the environment as emissions of nitrous oxide, nitric oxide, ammonia or molecular nitrogen to the atmosphere, or as nitrate or organic nitrogen to the hydrosphere before the food product is supplied to the consumer [26].

To assess the nitrogen intake ( $N_{cons}$ ) and food production N footprint ( $NF_{prod}$ ), we have used the data from [26]. The authors have provided NF factors calculated with the CAPRI model for individual crops and applicable to the quantity of food supplied at the farm gate. The data have been aggregated to the FBS crop groups, and  $NF_{prod}$  has been scaled to also include the losses of reactive nitrogen linked to wastages that occur between the farm gate and supply to the consumer. The quantity of these wastages is also given in the FBS.

The  $N_{cons}$  and  $NF_{prod}$  of avoidable food waste are calculated in analogy to the WF, as shown in equation (9) (table 2). Leip *et al* [26] also provide detailed data for the EU on the fate of the N surplus, and thus its potential contribution to adverse environmental impacts and associated damage costs [6, 7, 41–47]. Specifically, the shares emitted into the atmosphere as nitrous oxide ( $N_2O$ ), ammonia ( $NH_3$ ), nitrogen oxides ( $NO_x$ ) or dinitrogen ( $N_2$ ), or released to the hydrosphere via surface run-off or nitrogen leaching have been calculated on the basis of the life-cycle assessment model implemented in the CAPRI model for GHG accounting [48] and extended to cover nitrogen flows [26]. We have used their database of crop-specific emission unit flows to quantify the emissions

**Table 6.** Different components of the EU  $NF_{prod}$  in kg/cap/yr and Tg/yr (total 32.4 kg/cap/yr or 15.9 Tg yr<sup>-1</sup> for the EU population). The ‘nitrogen leaching and runoff’ component is dominant (36.4%), followed by the ‘N<sub>2</sub>’ component (29.3%). These values are comparable with earlier work [45].

	Emissions related to NFprod		Share of total NFprod
	kg/cap/yr	TgN/yr	%
N <sub>2</sub> O	0.58	0.28	1.8
NH <sub>3</sub>	6.22	3.03	19.2
NO <sub>x</sub>	0.19	0.09	0.6
N <sub>2</sub>	9.48	4.62	29.3
Nitrogen leaching and runoff	11.78	5.75	36.4
Nitrogen in animal waste and excess manure	4.02	1.96	12.5
Sum	32.35	15.78	100.0

of reactive nitrogen associated with food consumption in Europe.

## 3. Results


### 3.1. Values for FOOD\_CONS

As a first step, we calculated the total amount of food reaching consumers (FOOD\_CONS) based upon FAO FBS FOOD\_SUPPLY values for different product groups and the related water and NFs. These results are presented in table 5.

More particularly, table 5 shows:

- FOOD\_SUPPLY values for the EU for different product groups from the FAO FBS;
- FOOD\_CONS values after applying the correction factor CORR1 (equation (1)). All food reaching consumers (FOOD\_CONS) equals 789 kg/cap/yr;
- The FOOD\_CONS related  $WF_{cons,agr,gn}$  and  $WF_{cons,agr,bl}$  values (in l/cap/d). The total (sum of all product groups)  $WF_{cons,agr,gn}$  amounts to 3383 l/cap/d and the  $WF_{cons,agr,bl}$  to 270 l/cap/d. These values have already been presented in [9];
- The  $N_{cons}$  and  $NF_{prod}$  (in kg/cap/yr) values related to FOOD\_CONS. The total (sum of all product groups)  $N_{cons}$  amounts to 6.0 kg/cap/yr and the  $NF_{prod}$  to 32.4 kg/cap/yr.

The NF comprises  $N_{cons}$  and  $NF_{prod}$ . Total FOOD\_CONS  $NF_{prod}$  (N losses to the environment before the food product is supplied to the consumer) amount to 15.8 Tg N yr<sup>-1</sup>. About 30% of total N losses occur as molecular nitrogen ( $N_2$ ). They have no adverse effect on the environment and thus represent inefficient resource use [41] (see table 6 below). The ‘nitrogen leaching and runoff’ component is dominant (36.4%). These values are comparable with earlier work [45]. Up to 13% of the total losses



**Figure 2.** Distribution of total and avoidable EU consumer food waste in kg/cap/yr; statistical values in million tonnes per year (Mt/yr) and % of all food (789 kg/cap/yr or 385 Mt/yr) reaching consumers are also presented.

**Table 7.** Values (av = average; stdev = standard deviation, max = maximum; min = minimum) per product group of the green and blue  $WF_{\text{cons,agr}}$  of WASTE\_AV.


Product group	$WF_{\text{cons,agr}}$ of WASTE_AV (in l/cap/d)							
	Green				Blue			
	av	stdev	max	min	av	stdev	max	min
Cereals	57.603	24.768	97.730	16.850	3.589	1.543	6.090	1.050
Potatoes	8.349	4.110	14.850	1.650	1.518	0.747	2.700	0.300
Sugar	11.880	2.301	15.792	7.901	2.256	0.437	2.998	1.500
Pulses, nuts, oil crops	4.452	1.517	6.650	1.900	0.469	0.160	0.700	0.200
Crop oils	13.587	4.670	20.510	5.860	1.067	0.367	1.610	0.460
Vegetables	10.356	5.154	18.593	1.899	3.085	1.535	5.538	0.566
Fruit	17.131	8.354	31.235	4.646	4.469	2.179	8.148	1.212
Stimulants	24.018	4.621	32.178	16.103	0.149	0.029	0.200	0.100
Spices	0.675	0.128	0.900	0.450	0.075	0.014	0.100	0.050
Alcoholic beverages	4.168	0.987	5.880	2.521	0.298	0.071	0.420	0.180
Meat	105.609	52.666	191.846	25.449	6.945	3.463	12.615	1.673
Animal fats	4.930	1.684	7.210	2.060	0.287	0.098	0.420	0.120
Eggs	3.679	1.170	5.289	1.417	0.216	0.069	0.311	0.083
Milk (excl.butter)	10.557	5.065	28.437	3.840	0.825	0.396	2.222	0.300
Cheese	14.052	7.590	38.590	4.540	1.052	0.568	2.890	0.340
Cream	2.868	1.361	6.978	1.060	0.216	0.103	0.527	0.080
Fish	0	0	0	0	0	0	0	0

occur in manure and animal waste, for which data on purposeful reuse in the food chain were not available [26]. These flows have been included here as a conservative estimate, although it is likely that part of this N will be recovered and not dispersed in the environment.

### 3.2. Food waste

Figure 2 shows the calculated average amounts of food wasted by EU consumers, which consist of waste at the household and food service/catering sector levels. Total and avoidable waste amounts are given. The figure shows:

- Total waste averages 123 kg/cap/yr or 60 million tonnes (Mt) annually. This represents 16% of all food reaching consumers.
- Avoidable consumer food waste averages 97 kg/cap/yr or 47 Mt/yr. This represents 12% of all food reaching consumers.
- The range between minimum and maximum values of both total and avoidable food waste is very wide.
- Avoidable food waste represents by far the largest part of total food waste; non-avoidable food waste represents only a small fraction.



### 3.3. WF of avoidable food waste

The  $WF_{\text{cons,agr}}$  of avoidable food waste amounts from EU consumers are presented in figure 3 and table 7. Values per food product group and the total values are both presented. The total  $WF_{\text{cons,agr}}$  averages  $52 \text{ km}^3 \text{yr}^{-1}$  for green water and  $5 \text{ km}^3 \text{yr}^{-1}$  for blue water, in absolute volumes, or 294 l/cap/d for green water and 27 l/cap/d for blue water. There is a wide range between minimum and maximum  $WF_{\text{cons,agr}}$  amounts as the range in avoidable food waste amounts is large.

These average absolute volumes represent 8.2% of the EU green  $WF_{\text{cons,agr}}$  and 8.9% of the EU blue

$WF_{\text{cons,agr}}$ . They also represent 0.8% of the global green  $WF_{\text{cons,agr}}$  and 0.5% of the global blue  $WF_{\text{cons,agr}}$ .

The product group that accounts for the largest avoidable food waste  $WF_{\text{cons,agr}}$  is meat (average green  $18.8 \text{ km}^3 \text{yr}^{-1}$  or 106 l/cap/d and blue  $1.2 \text{ km}^3 \text{yr}^{-1}$  or 7 l/cap/d), as meat is a very water-intensive product [16, 28].

Other product groups with high amounts of avoidable food waste  $WF_{\text{cons,agr}}$  are cereals, cheese and crop oils. The blue  $WF_{\text{cons,agr}}$  of wasted fruit is the second largest after meat, i.e.  $0.8 \text{ km}^3 \text{yr}^{-1}$  or 4 l/cap/d, as much of the fruit which is either produced in and

**Table 8.** Values (av = average; stdev = standard deviation, max = maximum; min = minimum) per product group of the NF ( $N_{\text{cons}}$  and  $N_{\text{prod}}$ ) of WASTE\_AV.

Product group	NF of WASTE_AV (in kg/cap/yr)							
	$N_{\text{cons}}$				$N_{\text{prod}}$			
	av	stdev	max	min	av	stdev	max	min
Cereals	0.291	0.125	0.493	0.085	0.325	0.140	0.551	0.095
Potatoes	0.051	0.025	0.090	0.010	0.076	0.037	0.135	0.015
Sugar	0.000	0.000	0.000	0.000	0.023	0.004	0.030	0.015
Pulses, nuts, oil crops	0.009	0.003	0.014	0.004	0.005	0.002	0.007	0.002
Crop oils	0.000	0.000	0.000	0.000	0.019	0.006	0.028	0.008
Vegetables	0.044	0.022	0.079	0.008	0.066	0.033	0.119	0.012
Fruit	0.015	0.007	0.027	0.004	0.045	0.022	0.081	0.012
Stimulants	0.007	0.001	0.010	0.005	0.000	0.000	0.000	0.000
Spices	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Alcoholic beverages	0.005	0.001	0.007	0.003	0.060	0.014	0.084	0.036
Meat	0.144	0.072	0.262	0.035	1.550	0.773	2.815	0.373
Animal fats	0.000	0.000	0.000	0.000	0.024	0.008	0.035	0.010
Eggs	0.011	0.003	0.016	0.004	0.032	0.010	0.047	0.013
Milk (excl.butter)	0.027	0.013	0.074	0.010	0.143	0.069	0.385	0.052
Cheese	0.037	0.020	0.102	0.012	0.192	0.104	0.527	0.062
Cream	0.005	0.003	0.013	0.002	0.038	0.018	0.092	0.014
Fish	0.032	0.017	0.062	0.008	0.145	0.079	0.277	0.037

imported into the EU is irrigated [24]. Wasted vegetables and sugar also have a substantial blue  $WF_{\text{cons,agr}}$  component.

### 3.4. NF of avoidable food waste

The NF of avoidable food waste amounts from EU consumers is presented in figure 4 and table 8. Both the values per food product group and the total values are presented. The total  $N_{\text{cons}}$  averages 0.33 Tg N yr<sup>-1</sup> and the  $N_{\text{prod}}$  averages 1.34 Tg N yr<sup>-1</sup> in absolute quantities or 0.68 kg N/cap/yr and 2.74 kg N/cap/yr in per capita quantities. In addition, the range between minimum and maximum NF amounts is wide as the range in avoidable food waste amounts is large.

These average values represent 11.3% of the  $N_{\text{cons}}$  supplied to EU consumers and 8.5% of the EU  $N_{\text{prod}}$ .

The product group that accounts for the largest avoidable food waste NF is meat (average  $N_{\text{cons}}$  of 0.07 Tg N yr<sup>-1</sup> or 0.14 kg N/cap/yr and  $N_{\text{prod}}$  of 0.76 Tg N yr<sup>-1</sup> or 1.55 kg N/cap/yr).

Other product groups with high avoidable food waste NF values are cereals, cheese and milk.

## 4. Discussion


Our average total food waste estimate of 123 (min 55–max 190) kg/capita annually (kg/cap/yr) for EU consumers is about 20% higher than the estimated 101 kg/cap/yr total food waste (76 kg/cap/yr for households and 25 kg/cap/yr for the food service/catering sector) for the EU-27 by [10]. However, the latter value is well within the range we have defined.

We have estimated that almost 80% (97 (45–153) kg/cap/yr) is avoidable food waste. This average value

compares well with the FAO estimate of 95–115 kg/cap/yr of avoidable food waste for Europe [8], although the boundary conditions for our study are quite different to those in the FAO study. The FAO study only used UK avoidable food waste fractions, while our study used food waste fractions from six national studies. The UK presented the highest values among these countries. In our study, we computed consumer food waste from EU citizens in the 28 Member States, while the FAO study analysed the whole of Europe including Russia. In particular, in parts of Eastern Europe outside the EU the average amount of food reaching consumers (FOOD\_CONS) is lower compared to that inside the EU.

Our estimated ranges are quite wide because the six national estimates differ substantially, with the highest food waste fractions identified in the UK and the lowest in Romania. Data were gathered for both these studies by means of interviews/the keeping of food diaries. As regards the UK, this diary was complemented with: (1) detailed measurements of the weight and types of food and drink waste acquired from a list of consenting households; and (2) a synthesis of waste data from a list of local authorities. Indeed, household food waste is not easily quantified as it is disposed of either as: (1) municipal waste (solid and organic waste); (2) in the sewer; (3) home composting; or (4) feed for animals. Thus, municipal waste statistics underestimate total household food waste. This restricted the availability of suitable national data for EU Member States.

In addition, the aim of our study was also to differentiate between different product groups. As such, only six national studies qualified. For example, the study in Germany quantifies food waste amounts that



are similar to the average amounts in our studies. As shown in table 3, German food waste fractions (CORR2) amounts are average within the CORR2 range and, because of the country’s high population weighting, these values are quite dominant.

Nevertheless, even the lowest avoidable waste value within the range (45 kg/cap/yr) is substantial and shows the importance of food waste reduction. EU consumers are responsible for a large proportion of total food loss and waste along the whole food

supply chain, a large part of which is avoidable. A minimum of 45 kg/cap/yr equals about 125 g/cap/d—i.e. an average apple a day. A maximum of 153 kg/cap/yr equals about 420 g/cap/d—i.e. a small loaf of bread a day. Within our study, high total as well as avoidable waste amounts are observed for the cereals, fruit and vegetables product groups. A major reason for this is that these food groups have a relatively short shelf-life which means consumers often do not use them in time [29].



A large volume of green and blue water resources are associated with this avoidable waste, although the range in values is also wide. For instance, the average EU green  $WF_{\text{cons,agr}}$  of EU consumers' wasted food ( $52 \text{ km}^3 \text{ yr}^{-1}$ ) roughly equals the total green consumptive water used for crop production in Spain [16]. This means that all green water resources used annually for crop production in Spain—one of the EU's larger agricultural producers—are virtually thrown away by EU consumers through avoidable food waste. The average EU food waste blue  $WF_{\text{cons,agr}}$  ( $5 \text{ km}^3 \text{ yr}^{-1}$ ) is slightly higher than the total EU blue consumptive municipal water use ( $4 \text{ km}^3 \text{ yr}^{-1}$  or 23 l/cap/d) [24] or equals 1.5 times the annual flow of Italy's River Arno [48].

In addition, high NF values are related to these avoidable waste amounts, once again within a substantial range. The N contained in avoidable food waste averages 0.68 kg/cap/yr or almost 2 g/cap/d. The mean N footprint (any remaining N that was used in the food production process) is roughly  $1.34 \text{ Tg yr}^{-1}$ , which is equivalent to mineral fertiliser use in the UK and Germany combined [49].

Although the highest total and avoidable waste amounts are observed for the cereals, fruit and vegetables product groups, it is wasted meat that accounts for the highest related wasted water and nitrogen resources. For instance, the green  $WF_{\text{cons,agr}}$  of wasted meat is slightly higher than the total green consumptive water use for crop production in the UK ( $18.0 \text{ km}^3 \text{ yr}^{-1}$ ) [16]. The blue  $WF_{\text{cons,agr}}$  of wasted meat is roughly equal to the total blue consumptive municipal water use in Germany ( $0.6 \text{ km}^3 \text{ yr}^{-1}$ ) and France ( $0.6 \text{ km}^3 \text{ yr}^{-1}$ ) combined. This is because the production of livestock products, especially meat, is very resource intensive. It has already been observed by [9, 28, 50] that a reduction in meat intake in the average consumer's diet has the largest associated reduction effect on a consumer's  $WF_{\text{cons}}$ . This is important as it can show EU consumers that they can save a lot of resources by reducing their avoidable meat waste.

## 5. Concluding remarks

In our study, we have calculated both the total and avoidable food waste of EU consumers. We have also differentiated between different food product groups. Uncertainty is very high in these waste quantities, but has never been assessed in previous studies. Thus, for the first time, we have provided an estimate of the possible range, based upon statistical data from six EU Member States. Data from other Member States were either not suitable or did not exist. Our findings show that:

- Avoidable consumer food waste averages 97 kg/cap/yr, i.e. 12% of all food reaching consumers.
  - Avoidable food waste represents by far the largest part of total food waste.
  - Particular product groups are proportionally wasted more than others, especially vegetables, fruit and cereals. A major reason for this is that they have a relatively short shelf-life which means consumers often do not use them in time. Another reason is the fact that they are generally cheaper per weight unit as compared to other product groups like meat. Therefore consumers often tend to over-purchase the former.
  - There is a very wide range between minimum and maximum amounts of both total (from 55 to 190 kg/cap/yr) and avoidable (from 45 to 153 kg/cap/yr) food waste. This is due to the fact that lifestyles and purchasing power can differ substantially between Member States.
  - Even the lowest amount of avoidable food waste (45 kg/cap/yr) represents a substantial quantity. It is roughly equal to the weight of an apple a day or, for all the citizens in the EU together, 22 million tonnes of food each year. A maximum of 153 kg/cap/yr is equal to about 420 g/cap/d (i.e. a small loaf of bread a day) or for all the citizens in the EU together, 75 million tonnes of food each year. Consequently, reducing food waste should be a concern for every consumer.
- We have calculated water and nitrogen resources associated with avoidable food waste. To quantify the water resources related to these amounts of wasted food we have used the WF of consumption ( $WF_{\text{cons}}$ ) concept. In particular, we have assessed the green and blue  $WF_{\text{cons}}$  of agricultural products ( $WF_{\text{cons,agr}}$ ) for food wasted. To quantify the nitrogen resources related to these wasted amounts of food we have used the NF concept. In particular, we have assessed the nitrogen contained in food ( $N_{\text{cons}}$ ) and the food production NF ( $NF_{\text{prod}}$ ) for avoidable food waste. We have determined that:
- The average  $WF_{\text{cons,agr}}$  of avoidable food waste from EU consumers amounts to  $52 \text{ km}^3 \text{ yr}^{-1}$  green water and  $5 \text{ km}^3 \text{ yr}^{-1}$  blue water in absolute volumes or 294 l/cap/d green water and 27 l/cap/d blue water.
  - The average  $N_{\text{cons}}$  of avoidable food waste from EU consumers amounts to  $0.33 \text{ Tg N yr}^{-1}$  and the  $NF_{\text{prod}}$  to  $1.34 \text{ Tg N yr}^{-1}$  in absolute quantities or 0.68 kg N/cap/yr and 2.74 kg N/cap/yr in per capita quantities.
  - The range between minimum and maximum values of the  $WF_{\text{cons,agr}}$  and NF related to avoidable
- Total consumer food waste averages 123 kg/cap/yr, i.e. 16% of all food reaching consumers.



food waste is very wide because the range in food waste values is large.

- Among all the product groups, meat accounts for the largest avoidable food waste footprints ( $WF_{\text{cons, agr}}$  and NF), although wasted amounts are much smaller when compared to other food product groups. That is because meat production is very resource intensive. In other words, a small reduction in wasted meat already equates to a large reduction in wasted water and nitrogen resources. A similar observation was made by Cuellar and Webber [14] for energy. The authors state that wasted meat in the USA is responsible for the most wasted energy, despite a smaller relative fraction of meat being wasted compared with grains.
- This is also true for other livestock products such as milk and cheese, although to a lesser extent.
- On the other hand, wasted resources associated with fruit and vegetables are relatively low compared to other food product groups, although the wasted quantities from these product groups are high.

Theoretically, zero avoidable food waste is a possibility for EU consumers. This would not only save a lot of money for the consumers themselves, but also for local authorities which have to pay for food-waste collection and treatment. In addition, this would not only save a large volume of water and avoid losses of reactive nitrogen, but it would also preserve other natural resources such as phosphorus, land and energy. In a world with limited resources, food security can only be achieved by a more sustainable use of resources along with adaptations to our consumption behaviour, including the reduction or, ideally, the eradication of food waste [6, 7, 46].

## Acknowledgments

The authors would like to thank the following for their input and clarifications on food waste statistical data: Robert van Otterdijk of the FAO, Felicitas Scheider of BOKU in Vienna, and Violeta Stefan of Aarhus University.

## References

- [1] FAO, IFAD and WFP 2013 The multiple dimensions of food security *The State of Food Insecurity in the World 2013* (Rome: FAO)
- [2] Finucane M M et al 2011 National, regional, and global trends in body-mass index since 1980: systematic analysis of health examination surveys and epidemiological studies with 960 country-years and 9.1 million participants *Lancet* **377** 557–67
- [3] EUROSTAT 2014 on-line database (<http://epp.eurostat.ec.europa.eu>)
- [4] WHO 2014 *Global Health Observatory Data Repository* (<http://apps.who.int/gho/data/node.main?lang=en>)
- [5] Beddington J et al 2012 *Achieving Food Security in the Face of Climate Change: Final Report from the Commission on Sustainable Agriculture and Climate Change* (Copenhagen, Denmark: CGIAR Research Program on Climate Change, Agriculture and Food Security (CCAFS))
- [6] Foley J A et al 2011 Solutions for a cultivated planet *Nature* **478** 337–42
- [7] Godfray H C J et al 2010 Food security: the challenge of feeding 9 billion people *Science* **327** 812–8
- [8] Gustavsson J et al 2011 *Global Food Losses and Food Waste—Extent, Causes and Prevention* (Gothenburg, Rome: Swedish Institute for Food and Biotechnology (SIK), FAO)
- [9] Vanham D, Mekonnen M M and Hoekstra A Y 2013 The water footprint of the EU for different diets *Ecol. Indic.* **32** 1–8
- [10] Monier V et al 2010 Preparatory study on food waste across EU *27 Report for the European Commission [DG ENV—Directorate C] BIO Intelligence Service (BIOIS)*, Paris
- [11] Grizzetti B et al 2013 The contribution of food waste to global and European nitrogen pollution *Environ. Sci. Policy* **33** 186–95
- [12] WRAP 2008 *The Food We Waste* (Banbury: WRAP—Waste & Resources Action Programme)
- [13] Priefer C, Jörissen J and Bräutigam K-R 2013 Technology options for feeding 10 billion people—Options for cutting food waste, study for the European Parliament *Science and Technology Options Assessment (STOA)*
- [14] Cuellar A D and Webber M E 2010 Wasted food, wasted energy: the embedded energy in food waste in the United States *Environ. Sci. Technol.* **44** 6464–9
- [15] Hoekstra A Y and Wiedmann T O 2014 Humanity's unsustainable environmental footprint *Science* **344** 1114–7
- [16] Hoekstra A Y and Mekonnen M M 2012 The water footprint of humanity *Proc. Natl Acad. Sci. USA* **109** 3232–7
- [17] Leip A et al 2014 Nitrogen-neutrality: a step towards sustainability *Environ. Res. Lett.* **9** 115001
- [18] Leach A M et al 2012 A nitrogen footprint model to help consumers understand their role in nitrogen losses to the environment *Environ. Dev.* **1** 40–66
- [19] Weinzettel J et al 2013 Affluence drives the global displacement of land use *Glob. Environ. Change* **23** 433–8
- [20] Rööfs E et al 2013 Can carbon footprint serve as an indicator of the environmental impact of meat production? *Ecol. Indic.* **24** 573–81
- [21] Chapagain A and James K 2011 The water and carbon footprint of household food and drink waste in the UK *WWF and WRAP Report*
- [22] FAO 2013 *Food Wastage Footprint—Impacts on Natural Resources* (Rome: FAO)
- [23] Kummu M et al 2012 Lost food, wasted resources: global food supply chain losses and their impacts on freshwater, cropland, and fertiliser use *Sci. Total Environ.* **438** 477–89
- [24] Vanham D and Bidoglio G 2013 A review on the indicator water footprint for the EU28 *Ecol. Indic.* **26** 61–75
- [25] Vanham D and Bidoglio G 2014 The water footprint of agricultural products in European river basins *Environ. Res. Lett.* **9** 064007
- [26] Leip A et al 2014 The nitrogen footprint of food products in the European Union *J. Agric. Sci.* **152** 20–33 (supplement S1)
- [27] Food and Agriculture Organization 2013 *FAOSTAT On-line Database* (<http://faostat.fao.org>) (Rome: FAO)
- [28] Vanham D, Hoekstra A Y and Bidoglio G 2013 Potential water saving through changes in European diets *Environ. Int.* **61** 45–56
- [29] WRAP 2009 *Household Food and Drink Waste in the UK* (Banbury: WRAP—Waste & Resources Action Programme)
- [30] van Westerhoven M and Steenhuisen F 2010 *Bepaling Voedselverliezen Bij Huishoudens En Bedrijfs catering in Nederland* (Amsterdam: CREM)
- [31] Danish EPA 2012 *Kortlægning Af Dagrenovation I Enfamilieboliger* (Copenhagen: Miljøstyrelsen (Danish Environmental Protection Agency))
- [32] Silvennoinen K et al 2014 Food waste volume and composition in Finnish households *Br. Food J.* **116** 1058–68

- [33] Kranert M *et al* 2012 *Ermittlung der Weggeworfenen Lebensmittelmengen und Vorschläge zur Verminderung der Wegwerfrate bei Lebensmitteln in Deutschland* (Stuttgart: Institut für Siedlungswasserbau, Wassergüte- und Abfallwirtschaft, Universität Stuttgart)
- [34] Stefan V *et al* 2013 Avoiding food waste by romanian consumers: the importance of planning and shopping routines *Food Qual. Prefer.* **28** 375–81
- [35] Schneider F 2013 Review of food waste prevention on an international level *Proc. ICE—Waste and Resource Management* **166** 187–203
- [36] Rockström J *et al* 2009 Future water availability for global food production: the potential of green water for increasing resilience to global change *Water Resour. Res.* **45** W00A12
- [37] Hoekstra A Y *et al* 2011 *The Water Footprint Assessment Manual: Setting the Global Standard* (London, UK: Earthscan)
- [38] Thaler S *et al* 2012 Considerations on methodological challenges for water footprint calculations *Water Sci. Technol.* **65** 1258–64
- [39] Hoff H *et al* 2014 Water footprints of cities; indicators for sustainable consumption and production *Hydrol. Earth Syst. Sci.* **18** 213–26
- [40] Mekonnen M M and Hoekstra A Y 2011 National water footprint accounts: the green, blue and grey water footprint of production and consumption *Value of Water Research Report Series No. 50* UNESCO-IHE, Delft, The Netherlands
- [41] Pelletier N and Leip A 2014 Quantifying anthropogenic mobilization, flows (in product systems) and emissions of fixed nitrogen in process-based environmental life cycle assessment: rationale, methods and application to a life cycle inventory *Int. J. Life Cycle Assess.* **19** 166–73
- [42] Sobota D J *et al* 2015 Cost of reactive nitrogen release from human activities to the environment in the United States *Environ. Res. Lett.* **10** 025006
- [43] Van Grinsven H J M *et al* 2013 Costs and benefits of nitrogen for Europe and implications for mitigation *Environ. Sci. Technol.* **47** 3571–9
- [44] Weiss F and Leip A 2012 Greenhouse gas emissions from the EU livestock sector: a life cycle assessment carried out with the CAPRI model *agric Ecosyst. Environ.* **149** 124–34
- [45] Westhoek H *et al* 2014 Food choices, health and environment: effects of cutting Europe's meat and dairy intake *Glob. Environ. Change* **26** 196–205
- [46] Bodirsky B L *et al* 2014 Reactive nitrogen requirements to feed the world in 2050 and potential to mitigate nitrogen pollution *Nat. Commun.* **5** 3858
- [47] Sutton M A *et al* 2011 Too much of a good thing *Nature* **472** 159–61
- [48] Cortecci G *et al* 2009 Geochemistry of trace elements in surface waters of the arno river basin, northern Tuscany, Italy *Appl. Geochem.* **24** 1005–22
- [49] Gugele B *et al* 2014 *Annual European Union Greenhouse Gas Inventory 1990–2012 and Inventory Report 2014* European Commission (DG Clima) and EEA (European Environment Agency)
- [50] Vanham D 2013 The water footprint of Austria for different diets *Water Sci. Technol.* **67** 824–30